



RUNX1 gene

runt related transcription factor 1

Normal Function

The *RUNX1* gene provides instructions for making a protein called runt-related transcription factor 1 (RUNX1). Like other transcription factors, the RUNX1 protein attaches (binds) to specific regions of DNA and helps control the activity of particular genes. This protein interacts with another protein called core binding factor beta or CBF β (produced from the *CBFB* gene), which helps RUNX1 bind to DNA and prevents it from being broken down. Together, these proteins form one version of a complex known as core binding factor (CBF). The RUNX1 protein turns on (activates) genes that help control the development of blood cells (hematopoiesis). In particular, it plays an important role in development of hematopoietic stem cells, early blood cells that have the potential to develop into all types of mature blood cells such as white blood cells, red blood cells, and platelets.

Health Conditions Related to Genetic Changes

core binding factor acute myeloid leukemia

A rearrangement (translocation) of genetic material involving the *RUNX1* gene is found in approximately 7 percent of individuals with a form of blood cancer known as acute myeloid leukemia (AML). The translocation, written as t(8;21), combines genetic information from chromosome 21 and chromosome 8, fusing the *RUNX1* gene on chromosome 21 with a gene on chromosome 8 called *RUNX1T1* (also known as *ETO*). Because this genetic change affects CBF, the condition is classified as core binding factor AML (CBF-AML).

The resulting fusion protein, RUNX1-ETO, is able to form CBF and attach to DNA, like the normal RUNX1 protein; however, instead of turning genes on, it turns them off. This change in gene activity blocks the maturation (differentiation) of blood cells and leads to the production of abnormal, immature white blood cells called myeloid blasts. While t(8;21) is important for leukemia development, a mutation in one or more additional genes is typically needed for the myeloid blasts to develop into cancerous leukemia cells.

cytogenetically normal acute myeloid leukemia

rheumatoid arthritis


other disorders

Translocations and other types of mutations involving the *RUNX1* gene have been associated with different types of leukemia and related blood disorders, including acute lymphoblastic leukemia (ALL), chronic myelomonocytic leukemia (CMML), familial platelet disorder with predisposition to acute myeloid leukemia, and myelodysplastic syndromes (MDS). Depending on the type of mutation, these conditions can be related to impaired regulation of gene activity or loss of normal gene function. The *RUNX1* gene mutations associated with these diseases are somatic mutations and are not inherited. They are found only in certain cells of the body.

Chromosomal Location

Cytogenetic Location: 21q22.12, which is the long (q) arm of chromosome 21 at position 22.12

Molecular Location: base pairs 34,787,801 to 35,049,334 on chromosome 21 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- acute myeloid leukemia 1 protein
- AML1
- AMLCR1
- CBF-alpha-2
- CBFA2
- core-binding factor, runt domain, alpha subunit 2
- oncogene AML-1
- PEA2-alpha B
- PEBP2-alpha B
- PEBP2A2

- PEBP2aB
- polyomavirus enhancer-binding protein 2 alpha B subunit
- runt-related transcription factor 1
- RUNX1_HUMAN
- SL3-3 enhancer factor 1 alpha B subunit
- SL3/AKV core-binding factor alpha B subunit

Additional Information & Resources

Educational Resources

- The Cell: A Molecular Approach (second edition, 2000): Structure and Function of Transcriptional Activators
https://www.ncbi.nlm.nih.gov/books/NBK9904/#_A1003_

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28RUNX1%5BTI%5D%29+OR+%28runt-related+transcription+factor+1%5BTI%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5BIa%5D+AND+human%5Bmh%5D+AND+%22last+360+days%22%5Bdp%5D>

OMIM

- PLATELET DISORDER, FAMILIAL, WITH ASSOCIATED MYELOID MALIGNANCY
<http://omim.org/entry/601399>
- RUNT-RELATED TRANSCRIPTION FACTOR 1
<http://omim.org/entry/151385>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
<http://atlasgeneticsoncology.org/Genes/AML1ID52.html>
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=RUNX1%5Bgene%5D>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=10471

- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/861>
- UniProt
<http://www.uniprot.org/uniprot/Q01196>

Sources for This Summary

- Goyama S, Mulloy JC. Molecular pathogenesis of core binding factor leukemia: current knowledge and future prospects. *Int J Hematol*. 2011 Aug;94(2):126-33. doi: 10.1007/s12185-011-0858-z. Epub 2011 May 3. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21537931>
- Goyama S, Schibler J, Cunningham L, Zhang Y, Rao Y, Nishimoto N, Nakagawa M, Olsson A, Wunderlich M, Link KA, Mizukawa B, Grimes HL, Kurokawa M, Liu PP, Huang G, Mulloy JC. Transcription factor RUNX1 promotes survival of acute myeloid leukemia cells. *J Clin Invest*. 2013 Sep;123(9):3876-88. doi: 10.1172/JCI68557. Epub 2013 Aug 27. Erratum in: *J Clin Invest*. 2013 Nov 1;123(11):4979.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/23979164>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3754260/>
- Huang G, Shigesada K, Ito K, Wee HJ, Yokomizo T, Ito Y. Dimerization with PEBP2beta protects RUNX1/AML1 from ubiquitin-proteasome-mediated degradation. *EMBO J*. 2001 Feb 15;20(4):723-33.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11179217>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC145428/>
- Lam K, Zhang DE. RUNX1 and RUNX1-ETO: roles in hematopoiesis and leukemogenesis. *Front Biosci (Landmark Ed)*. 2012 Jan 1;17:1120-39. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22201794>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3433167/>
- OMIM: RUNT-RELATED TRANSCRIPTION FACTOR 1
<http://omim.org/entry/151385>
- Ran D, Shia WJ, Lo MC, Fan JB, Knorr DA, Ferrell PI, Ye Z, Yan M, Cheng L, Kaufman DS, Zhang DE. RUNX1a enhances hematopoietic lineage commitment from human embryonic stem cells and inducible pluripotent stem cells. *Blood*. 2013 Apr 11;121(15):2882-90. doi: 10.1182/blood-2012-08-451641. Epub 2013 Jan 31.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/23372166>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3624936/>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/RUNX1>

Reviewed: November 2013
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services